


International Journal of Agricultural Science & Technology




This Journal is an academic
and peer-reviewed publication
(Print ISSN 2319-880X)

An International Refereed Journal



ARP. Give your IDEAS an Expression

www.manishanrpp.com



Cite this volume as
4(1)IJAST(2015)
and so on....

ABOUT THE JOURNAL

INTERNATIONAL JOURNAL OF AGRICULTURAL SCIENCE AND TECHNOLOGY is a biannual academic and peer-reviewed Journal published by **ACADEMIC AND RESEARCH PUBLICATIONS in Collaboration with JPMS Society**. JPMS Society is a Society registered under the Societies Registration Act and its Registration No. is 1649/1986-87. It was published from year i.e. 2012. The ISSN of the JOURNAL is **2319-880X**.

OBJECTIVE OF THE JOURNAL

To promote and encourage specially YOUNG SCIENTISTS to take active part in research and get acquainted with the latest development and research in the field of environmental science and technology. To promote cooperation in the pursuit of research in general and to exchange and contribute to the progress in the field of Agricultural Science and technology in particular.

CALL FOR PAPERS

We invite you to submit high quality papers for review and possible publication in all areas of Agricultural Science and technology which includes Soil Science, Agri. Entomology, and Gnd.water Pollution, Limnology, Toxicology, Animal Husbandry, Biological and Chemical Control, Seed Science and Technology, Agricultural Waste, Any other related topics. All authors must agree on the content of the manuscript and its submission for publication in this Journal before it is submitted to us. Manuscripts should be submitted by e-mail to the Editor at **manisha_npp@yahoo.com**.

COVERAGE OF THE JOURNAL

- | | |
|-------------------------------------|-------------------------------|
| 1. Soil Science | 2. Water & Soil Pollution |
| 3. Agricultural Entomology | 4. Aquaculture |
| 5. Groundwater Pollution | 6. Pesticides |
| 7. Limnology | 8. Sericultural Biotechnology |
| 9. Toxicology | 10. Plant Protection Science |
| 11 Animal Husbandry. | 12. Meteorology |
| 13. Biological and Chemical Control | 14. Agronomical Science |
| 15. Seed Science and Technology | 16. Forest and Forestry |
| 17. Agricultural Waste | 18. Any other related topics |

TYPES OF PAPERS ARE INVITED

Following types of papers are invited for publication in this Journal :-

- | | |
|--|------------------|
| a) Original Research Papers of Scientific values | b) Review Papers |
| c) Short Communications | d) Case Reports |
| e) Letters to the Editor | f) As you see |

REVIEWERS PROCESS

All manuscripts are reviewed by an editor and members of the Editorial Board or qualified outside REVIEWERS. Decisions will be made as rapidly as possible and the Journal strive to return reviewer's comments to authors within 6 weeks. The Editorial Board will re-review manuscripts that are accepted pending reviewers. It is the goal of the this Journal to publish manuscripts within 4 weeks after submission after getting O K report from the Author.

CONTACT US

For quick reply, please note change of address and contact them directly by Post or email:-

For publication of your article, Acceptance letter, Review Reports, Status Report, and all other queries related to your articles, should be sent directly to the Editor-in-Chief, whose address is as follows:

Prof. Manik Sinha, THE EDITOR-IN-CHIEF

Email : **manik.sinha2@gmail.com**, Contact at : 09415155631

For all publication matters related to the Journals Acceptance letter for publication of articles, Invoice, Reprints etc. should be sent directly to the PUBLICATION EDITOR whose address is as follows :
To,

Ex. Manisha Verma, B. Sc., B. Tech.

Publication Editor (Chief Executive Director)

Academic And Research Publications

H. Office: EC 41, Maya Enclave, New Delhi -110064

Email : **manisha_npp@yahoo.com** **www.manishanpp.com**

IJAST

....

International Journal Of Agricultural Science & Technology

Volume 4, Issue 1

February 2015

Cite this volume as 4(1)IJAST(2015) and so on....

“ This journal is Indexed/abstracted in Indian Science Abstract ”

© Journal on Agricultural Science and Technology. All rights reserved. No portion of material can be reproduced in part or full without the prior permission of the Editor.

Note : The views expressed herein are the opinions of contributors and do not reflect the stated policies of the

ACADEMIC AND RESEARCH PUBLICATIONS AND JPMS SOCIETY

This Journal is an academic and peer-reviewed publication (Print ISSN 2319-880X)



An International Refereed Journal



ARP. Give your IDEAS an Expression

www.manishanpp.com



Academic And Research Publications

H. Office: EC 41, Maya Enclave, New Delhi -110064

International Journal Of Agricultural Science & Technology

Volume 4, 2014

February 2015

Issue 1, 2015

Editorial Board

Editor-in-Chief

Prof. Manik Sinha

Former Dean, Faculty of Law,
Dr R.M.L Awadh University, Faizabad (UP),
Senior Advocate, Govt Of India, High Court, Lucknow
Email: manik.sinha2@gmail.com

Chief Editor

Prof. Padmakar Tripathy,
Dept of Meteorology,
N D University of Agri. and Tech.,
FAIZABAD - 224 229 U P
E-mail: pmtmausam@gmail.com

Associate Editor

Dr. K. K. Singh,

Head, Agromet Services &
Scientist-F
India Meteorological Department
Lodhi Road, New Delhi - 110003

Dr. A. P. Dubey, Professor
(Agromet)

Division of Agrometeorology
C.S.A. U of A & , Kanpur-208002
Dr. S. R. Mishra Asstt. Professor
Deptt. of Agril. Met.
N.D.U.A. & T., Faizabad-224229
e-mail: srmmausam@india.com

Dr. A. K. Singh Asstt. Professor
Department of Crop Physiology
N. D. U A. & T,
FAIZABAD - 224 229
E: assinghkumar3@gmail.com

Associate Editors

Mrs. Chandra Kanti Sinha
Retd. Lecturer in Botany(Principal Grd.),
Government College Faizabad U.P.
Email: chandrakantisinha@yahoo.com

Assistant Editors

Dr. Prateek Tripathi

Molecular & Computational Biology
University of Southern California
TRF 108, 3430 S Vermont Ave
Los Angeles CA-90089
Email ; tprateek@dornsife.usc.edu

Dr. Vishva Deepak Tripathi

Research Associate
NRC for Agroforestry,
Near Pahuj Dam
Gwalior Road, Jhansi -284 003, U.P.

Dr. S. B. Yadav

Department of Agromet
A A U, Anand, Gujarat,
e-mail: sbmet@yahoo.in

Dr. R. P. Kaushal Asstt. Prof.

Deptt. of Statistics & Social Science
MSKJUA&T, Banda - 210001 (UP)
Email: kaushal_rajendra@india.com

Executive Editor

Prof S. K. Tripathi

Department of Water Resources De-
velopment and Management
I.I.T., Roorki-247667,
e-mail: sankufwt@gmail.com
sankufwt@iitr.ernet.in

Coordinating Editor

Dr. Arvind Kumar Srivastava,

Dept of Meteorology,
N D University of Agri. and
Tech., Faizabad, U P.
E: arvindmausam@gmail.com

Technical Editor

Dr. A. K. Srivastava, Principal
Scientist (Agromet)

I I S R Lucknow
E: arunkumarsrivastava2@gmail.com

Publication Editor

Dr. Manisha Verma, B.Sc., B.Tech.

Publication Editor (Chief Executive Director)

Academic And Research Publications

H. Office: EC 41, Maya Enclave, New Delhi -110064

Email : manisha_npp@yahoo.com www.manishanpp.com

© Journal on Agricultural Science and Technology. All rights reserved. No portion of material can be reproduced in part or full without the prior permission of the Editor.

Note : The views expressed herein are the opinions of contributors and do not reflect the stated policies of the Academic And Research Publications. Correspondence: All enquiries, editorial, business and any other, may be addressed to: The Editor-in-chief, International Journal of Agricultural Science and Technology (IJAgST), H. Office: EC 41, Maya Enclave, New Delhi -110064 (INDIA)

Email : manik.sinha@ymail.com; manisha_npp@yahoo.com, www.manishanpp.com.

International Journal Of Agricultural Science & Technology

Members of Editorial Board

Dr. V. N. Pathak

Asstt. Prof., Deptt of GPB
S.M.M. Town College, Ballia, U.P.
E: vijayanand.pathak76@gmail.com

Dr. Alok Kumar Singh

Deptt of Crop Physiology
N. D. U A. & T, FAIZABAD-224 229
.e-mail: aloksingh.agri@gmail.com

Dr. B. P. Shahi

Asstt Prof./SMS (Horticulture)

Krishi Vigyan Kendra, Meerut.
e-mail: bhshahi1975@yahoo.com

Dr. P. K. Singh

India Meteorological Department
Mausam Bhavan, Lodhi Road
New Delhi-110003, India.
e-mail: pksingh66@gmail.com

Prof J. P. Srivastava

Head, Deptt of Plant Physiology
Institute of Agricultural Sciences
Banaras Hindu University, Varanasi,

Dr. Vyas Pandey

Prof. & Head, Deptt of Agromet
A A U, Anand, Gujarat,
e-mail: pandey04@yahoo.com

Dr. Akhilesh Srivastava

Assoc. Prof., Horticulture
M P U of Agril. & Tech., Udaipur,

Dr. Anil Kumar Singh

Asstt. Prof., Agronomy
N.D.U.A.& T., Faizabad.
e-mail: aksmausam@gmail.com

Dr. Neeraj Kumar

Astt Prof, Deptt of Soil Science,
N. D. U A. & T, FAIZABAD - 224 229
e-mail: nksoilchem@gmail.com

Copyright: Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

International Journal Of Agricultural Science & Technology

Volume No. 4

Feb 2015

Issue No. 1, 2015

C o n t e n t s

S. No.	Title	Page No.
--------	-------	----------

- | | | |
|----|---|----|
| 1. | Characteristic of Fog, Foggy Weather and its Impact on Agriculture. | 01 |
|----|---|----|

Anil Kumar, Raj Singh and S. Singh

- | | | |
|----|--|----|
| 2. | Economical Analysis Of Potato Production: Nature And Extent Of Marketable & Marketed Surplus | 05 |
|----|--|----|

Pramod Kumar, R. P. Kaushal, U. A. Siddiqui and Daya Ram

- | | | |
|----|--|----|
| 3. | Effect Of Profenofos (Insecticide) on Hematological Parameters of Occupationally Exposed Workers | 16 |
|----|--|----|

Sadhana Kesharwani And Khushboo Sahu

- | | | |
|----|--|----|
| 4. | Diversity Of Macrophyte Community Of A Lentic Water Body Of Jabalpur | 23 |
|----|--|----|

Dr. Reeta Solanki



© Journal on Agricultural Science and Technology. All rights reserved. No portion of material can be reproduced in part or full without the prior permission of the Editor.



CHARACTERISTIC OF FOG, FOGGY WEATHER AND ITS IMPACT ON AGRICULTURE

Anil Kumar, Raj Singh and S. Singh

Department of Agricultural Meteorology
COA, CCS HAU, Hisar-125004

Email : anilmeteo@gmail.com

(Date of Receipt :14-12-2014 ;

Date of Acceptance for Publication : 17-01-2015)

Abstract

This work is on the basis of different literature survey which is base on the real time observation are observed at during the foggy weather condition at Research Farm, Department of Agricultural Meteorology, CCS HAU, Hisar during the winter season of subsequent year on rabi crop, several significant impact to promoting the fungal diseases if prolonged days of foggy weather also minimized the dispersion rate of beneficial insect over the crop and their ideal time of pollination by beneficial insect will also change, Insecticide and liquid fertilizer over the crop field are not recommending spraying during dense or thicker fog condition, adversely affect due to reduced solar radiation as PAR which is influencing for photosynthesis and reduces the evapotranspiration and photosynthesis rate of different field crop.

Key words: Fog, Mist, Foggy Weather, Impact on Rabi Crop, Pest Dispersion.

Pages :04

References: 07

INTRODUCTION

Brief About Fog

The particular weather condition would occur in formation of fog in the environment and lot of moisture level, average relative humidity level was higher. The regular phenomena in the environment during the winter seasons (India's most of the part is covered by fog during the winter season) in regions like North

West, North East and Central India but few days with less thicker fog observed in the Western Part of India also. Fog consist small tiny suspended droplets of water molecules, which may be settling out and evaporating, which restrict the clear horizontal visibility and persist for some period of time during early morning and night time, generally sun rise with high intensity insolation started its quickly dissipated. Fog is generally a collection of liquid water



droplets suspended in air at or near the Earth's surface, Fog can be considered a type of low-lying cloud, and is heavily influenced by nearby bodies of water, topography, wind conditions, and even human activities. In turn, fog has affected many human activities, such as shipping and transport, warfare, and culture. Fog can contribute significantly to crop water needs and can be measured by covering the funnel of a rain gauge with a set of fine wires. Quantitative data on fog precipitation may not be available. However, nomograms for predicting occurrence of fog at airports are available with forecasters and the same can be adopted for use in agricultural weather forecasts. The requirement of meteorological conditions for fog formation are the same as for dew those formation except for the need for absence of air-turbulence or current (vertical movement of air) in the air layers close to the ground and crop-canopy temperature being lower than the screen temperatures. Fog season is generally considered from November to February, with the months of December and January witnessing the longest durations of dense fog in North-west plains of India Singh and Singh 2010; Verma, 1989; Samui and Gupta 1994).

Depth of Fog

Fog may vary in depth occurring within the lowest meter of the atmosphere and extending up to 1000m in height. Occasionally the fog layer may be somewhat elevated above the surface, particularly during the fog dispersion, breakup process. For condensation, the relative humidities of the air do not necessarily need to be 100% and in fact may be as low as

80–90% both during and after formation. These conditions are mostly associated with a temperature dew point spread of approximately 3 °C and a stable layer of atmosphere.

Fog or Foggy Weather

During the foggy weather conditions the early morning have same mercury reading in Dry bulb temperature and wet bulb temperature, there is less possibility to mix the more amount of water vapour in the surrounding atmospheric air and relative humidity value becomes 100 percent. It creates the situation of different parameters value become equal like saturated vapour pressure (SVP), actual vapour pressure (AVP), dew point temperature, dry bulb and wet bulb temperature during early morning (at 0730 hrs) and then dissipated with start the insolation reaching on the earth surface.

MATERIALS AND METHODS

This work is on the basis of different literature survey and real time observation which were observed at real time during the foggy weather condition at Research Farm, Department of Agricultural Meteorology, CCS HAU, Hisar during the winter season of subsequent year on rabi crop. Different types of observation also monitoring for the pest dispersion and its dynamic during the period of Fog/mist or foggy weather condition.

Observation of Fog

Now-a-days various methods of taken observation and the basis of their uses in the site specific like agrometeorological ob-



server, breeder, insect ecologist, entomologist and plant pathologist for their own research purpose. Various methods and instrumentation used to determine visibility are presented in the visibility portion, but most of the site there is necked eyes observation on the visibility basis, visibility will be decided by human being eye observation in most of the agrometeorological observatory and then interpreted by the meteorological observer on the basis of their visibility from the standing point to clear distinct vision.

Types of observation

The most obvious types of observations of fog are its occurrence, reduction of visibility and fog's thickness or intensity, color (of limited use in reporting), duration, and extent. The traditional method of observation is on the basis of human eye observation, how far one can clearly see. The two basic processes responsible for formation, as well as duration, are radiation and advection which includes vertical mixing of air). It is based upon these observations that fog is often referred to as 'a cloud on the ground' and which consists of visible hydrometeors.

RESULTS AND DISCUSSION

The several significant advantages of fog/mist or foggy weather over the Agriculture during the winter season in north west, north east part of country. The surface grass hopper activity will minimized in the wheat crop field, Minimum termite infestation in cultivate field crop (i.e. wheat), the average higher humidity level prevailed during foggy weather to prevent the frosting condition somehow in Mustard vegetable crops and small horticultural

crop, Dispersion rate of insect and pest over crop will minimized. During the dense fog weather condition there will be chance to receive only diffuse or sky radiation. The minimum temperature was value observed over the normal and significant fall in the day maximum temperature could be occurs at the place there dense fog or mist weather was occurred during winter season. Dis-advantage of fog or foggy weather over the Agriculture, Due to high humidity level in foggy weather condition, possibility to occurrence of white rust and alternaria blight in the mustard crop. Pest and fungal diseases infestation would be maximized mist/foggy weather condition, increase the population of sucking pest due to higher humidity level which damage the tender part of crop. Pollination would be minimized due to higher, humidity level in self or cross pollination field crop, vegetable etc or Impact vegetables and fruits adversely if foggy and overcast conditions continue. and reduced the evapotranspiration and photosynthesis rate of different crops. Promote skin disease/infection of Animal/cattle. To minimized assimilation rate or CHO group accumulation or carbohydrate over crop or plant due to less PAR insolation/absorption. Dense fog and mist continue to occur for some more days then it can cause diseases like late blight on potato crop. The movement of pollinator or beneficial insect will also minimized due to high relative humidity present in the environment or surrounding lower atmospheric boundary layer. Using the weedicide, fungicide, Insecticide and liquid fertilizer over the crop field are not recommending spraying during dense or thicker fog condition due to leaf wetness duration will be increase to permit the fungal disease.



CONCLUSION

The foggy weather is somehow advantageous to the agriculture crop as it meet the some requirement of irrigation, weedicide, fungicide, Insecticide and liquid fertilizer over the crop field are not recommended to spraying during dense or thicker fog condition due to the prolong wetting time of crop leaves, reduces the dispersion rate of different insect pest and promote the fungal diseases of crop, adversely affected due to reduced the solar radiation as PAR which is influencing for photosynthesis and reduces the evapotranspiration and photosynthesis rate of different field crop, the infestation of insect will also become lower along with dispersion over the crop canopy.

REFERENCES

1. Croft, P. J. **(2003)**. Fog. University of Louisiana at Monroe, Monroe, LA, USA.
2. <http://en.wikipedia.org/wiki/Fog>.
3. http://www.agrometeorology.org/files-folder/repository/gamp_chapt4.pdf.
4. Samui, R.P. and Gupta, D. C. **(1994)**. Fog in relation to elevation and topographical features at two stations in Sikkim. *Mausam*, Vol. 45 (4), pp. 369-371, 1994.
5. Anil Kumar and V. Pande. **(2012)**. The effect of weather parameters, cultural practices on growth and yield of Soybean (*Glycine Max. [L] (Merrill)*) under different date of sowing and four levels of spacing. *International Journal of Agricultural Science & Technology*, Vol.1(1),pp. 49-60.
6. Singh, S. and Singh, D. **(2010)**. 5th International Conference on Fog, Fog Collection and Dew Münster, Germany, 25–30 July 2010 FOGDEW2010-100.
7. Verma, R.K. **(1989)**. Fog over Lucknow airport. *Vayu Mandal*, 19 (1-2): 39-42, 1989.





ECONOMICAL ANALYSIS OF POTATO PRODUCTION: NATURE AND EXTENT OF MARKETABLE & MARKETING SURPLUS

Pramod Kumar¹, R. P. Kaushal², U. A. Siddiqui³ and Daya Ram⁴

¹Assistant Professor, C.C.S.M. BardariBarabanki, U.P., (India)

²Assistant Professor, College of Agriculture, MSKJUAT, Banda

³ Assistant Professor, Agriculture Extension, Chandra Bhanu Gupta Agricultural P.G. College, Lucknow

⁴ Associate Professor, Agriculture Extension, Central Agricultural University Imphal

Email : kaushal_rajendra@india.com

(Date of Receipt :07-01-2015 ;

Date of Acceptance for Publication : 16-02-2015)

Abstract

The present study was conducted in Jaunpur district of Uttar Pradesh using the Multi-Stage Research Design. Total 100 respondents (44 marginal, 26 small, 18 medium and 12 large) were selected based on proportional random sampling method. The data were collected using structured interview schedule. The data collected were then analyzed using appropriate statistical tools namely, frequency, percentage, mean, and standard deviation. It may be concluded that overall, cost of cultivation per hectare was observed Rs. 31873.42 the major share of cost was 41.32 per cent of seed followed by 22.75 per cent family labour, 8.20 per cent manure and fertilizer respectively. On an average gross return, net return, family labour income, farm investment income and farm business income Rs. 98965.25, Rs. 60424.02, Rs. 67676.25 and Rs. 44859.12 respectively. Overall cost of production per quintal on the basis of cost C1 and C was found Rs. 157.91 and Rs. 175.45 respectively. On an overall average, input-output ratio was found 2.67 per cent. Overall average of marketable surplus came to 37.95 quintal i.e. 82.53 per cent of the total production and overall average of marketed surplus was recorded to 36.56 quintal per farm i.e. 79.51 per cent of the total production.

Key Words: Potato Production, Cost, Marketing and Measures.

Pages: 11

References: 25

INTRODUCTION

The original home of Potato is the Andean Plateau of South America. It was introduced into Europe in the second



half of the 16th Century but gained recognition as an inexpensive and a nutritive food only during the 18th Century. Potato is the most abundantly produced edible food in the world next to cereal. Potato is a wholesome food. Apart from starch of which it is a rich source, it provides essential body-building substances such as starch, vitamins, minerals and protein. Potato is one of the richest sources of calories needed to maintain day-to-day output of human energy. Seventy five per cent of the total Indian population depends on agriculture. The potato crop is particularly suited to Indian conditions. Besides ecological and agronomic factors, economic and social consideration also favors a rapid expansion for its cultivation. Potato enjoys a wide range of seasonal adaptability. The Gangetic plains of Uttar Pradesh contributes annually about 35 per cent of the total area under potato cultivation in the country. Here, the sowing time of the crop can be extended over a period of three months, from October to the end of December, an advantage which perhaps no other crop enjoys. In Southern India, where summer temperatures in the plateau region are somewhat milder, two crops, one in winter and the other in summer, can be raised. It should be recognized that very few crops can be raised successfully both in the rabi and the kharif seasons in the same track. In the Nilgiri Hills, three potato crops are raised almost in succession, the planting month's beings April, August, and January (Anonymous 2008).

Wide flexibility in planting time of a semi-perishable commodity like potato is of great significance, since it makes efficient planning and adjustments in

cropping sequence possible. Almost all crops grow, in one season or the other, in the plains of the country but only a few of them, of which potato is the chief one, grow successfully at high altitudes. In a country with limited resources, where the nutritional level of the population has to be maintained under inhospitable conditions, potato has a special place for culture. Potato can be grown almost on any type of soil, except alkaline soil, though it is tolerant even to saline conditions. It thrives best in sandy or sandy loam soils. A cereals crop cannot be grown successfully in poor types of loose, shifting soils. But unlike cereals, potato builds up food materials underground and thrives in a loose soil which allows its root system to spread and the young developing underground tubers to grow freely. It is, thus possible to get excellent potato crops even under a river-bed system of cultivations, as at Ahmedabad in Gujarat State. On the other hand, potato is by no means allergic to heavier types of soils suitable for rice or jute cultivation. Rice-potato and jute-potato are established rotations practiced in West Bengal and some parts of Eastern Uttar Pradesh.

Potato is one of the most important food crop and fourth in total production among food crops in the world. India is the 3rd largest potato producer in the world. The production of potato in the world is approximately 311.4 million tonnes with an area under potato 19.22 million hectare. An average productivity of potato is 16.20 tonnes/hectare, while, in India, production of potato was about 24 million tonnes with an area of 1.2 million hectares and productivity of 19.7 tonnes/hectare (Singh and Kumar, 2004).

The main potato producing states, in order of importance are Uttar Pradesh, Bihar, West Bengal, Assam, Punjab, Madhya Pradesh, Himachal Pradesh, Haryana, Maharashtra, Karnataka, Meghalaya and Tamil Nadu. In U.P., potato is cultivated in about 0.432 million hectares with the production of about 7.68 million tonnes. The productivity of potato in U.P. is 17.78 tonnes/hectare (Singh and Kumar, 2004).

Potato is one of the most important cash crop of district Jaunpur (U.P.) and has significant importance in the economy of the district. Area under potato in this district during 2001-02 was 0.117 lakh hectare while total production was 2.56 lakh metric tonnes. The productivity of potato in Jaunpur district was 21.82 tonnes /hectare but so far, not any systematic study on economics or potato production has been conducted. Keeping in view the above-mentioned facts, to study about costs and returns of potato cultivation and to work out the marketable and marketed surplus of potato.

METHODOLOGY

Multi-stage purposive cum-random sampling technique was used to selected district, block, villages and respondents. Jaunpur district of Eastern U.P. was selected purposively, because of higher concentration of area and production of potato in this district. Out of 19 blocks, one block namely, Shahganj was selected for the present study because of higher concentration of potato production in this block as compared to other blocks. A list of all villages of Shahganj block was prepared according to acreage under potato cultivation. Two nucleus villages were selected (one near city and another 20 km

away from city) to farm clusters with four surrounding villages of each nucleus villages. Thus, 10 villages were selected for the present study. A list of all the potato growers of 16 selected villages were prepared and classified into marginal (below 1 ha), small (1-2 ha), medium (2-3 ha) and large (3 & above ha). Further 10 farmers from each selected villages were randomly selected in proportion to their size groups of farm selected. Total 100 respondents (44 marginal, 26 small, 18 medium and 12 large) were selected based on proportional random sampling method. The data were collected using structured interview schedule. The data collected were then analyzed using appropriate statistical tools namely, frequency, percentage, mean and standard deviation.

$$\text{Weighted mean} = \frac{\sum WX_i}{\sum W}$$

Where,

W = Weight of X_i

X_i = Variable

RESULTS AND DISCUSSIONS

Cost of Cultivation of Potato

Table 1 revealed that on an average, cost of cultivation of potato per hectare was observed Rs. 32673.64 on first cluster. Average cost was found maximum on marginal farms i.e. Rs. 33108.71 followed by small farms Rs. 32958.63, medium farms Rs. 31945.93 and large farms Rs. 31552.43. Higher cost on marginal farms was mainly associated due to comparatively more expenditure on family labour. Whereas in case of second cluster, on an average, cost of cultivation were per hectare observed Rs. 31073.19. Average cost was higher on marginal farms i.e. Rs. 31920.63 followed by small farms Rs. 30904.21, medium farms Rs. 30172.73 and large

farms Rs. 29682.71. Higher cost on marginal farms was mainly due to comparatively more expenditure on family labour. The study further reveals that maximum cost was incurred on seed i.e. 42.29 per cent, followed by family labour 21.65 per cent, manure and fertilizers 8.20 per cent, irrigation charges 3.86 per cent, hired labour 3.34 per cent, tractor and machinery Charges 2.82 per cent, bullock labour 2.56 per cent and plant protection 1.71 per cent, respectively on first cluster farms.

The costs incurred on other items like interest on working capital and rental value of owned land were almost the same. The maximum share among these items was rental value of owned land being 11.63 per cent of the total cost per hectare.

Whereas in case of second cluster, it was 40.28 per cent on seed cost, followed by family labour 23.92 per cent, manure and fertilizers 8.20 per cent, irrigation charges 4.41 per cent, hired labour 3.29 per cent, tractor and machinery charges 2.86 per cent, bullock labour 2.34 per cent and plant protection 1.53 per cent, irrespective of size of sample farms. The costs incurred on other items like interest on working capital and rental value of owned land were almost the same. The maximum share among these items was of rental value of owned land being 11.26 per cent of the total cost per hectare.

The study further reveals that overall average cost per hectare was observed Rs. 31873.42 and major components of cost was 41.32 per cent on seed cost followed by family labour 22.75 per cent, manure and fertilizers 8.20 per cent, irrigation charges 4.13 per cent, hired labour 3.31 per cent, tractor and machinery charges 2.84 per cent, bullock labour 2.45 per cent and plant protection 1.62 per cent.

The cost incurred on other items like interest on working capital and rental values of owned land were almost the same. The maximum share among these items was of rental value of owned land being 11.45 per cent of the total cost per hectare.

It can be inferred from the Table 1 that, average cost (per hectare) in second cluster was more as compared to first cluster due to employment of more family force on second cluster.

Measures of Costs and Returns of Potato

Costs and returns of potato under different size group of sample farms are presented in Table 2. The cost C1 was accounted for Rs. 35241.24, Rs. 33303.60, Rs. 30580.53 and Rs. 29282.21 on marginal, small, medium and large size of sample farms respectively with an average of Rs. 33193.44 for first cluster whereas, it was Rs. 35413.68, Rs. 31611.90, Rs. 30192.41 and Rs. 28027.87 on marginal, small, medium and large size of sample farms respectively with an average of Rs. 32599.09 for second cluster. Overall average of cost C1 was observed Rs. 32891.27 for both the clusters. The findings were in line with the research results of Keruret al. (1998)

Similarly, cost C., was worked out i.e. Rs. 39041.23, Rs. 37103.60, Rs. 34380.53 and 33082.21 on marginal, small, medium and large size of sample farms respectively with an average of Rs. 36983.44 for first cluster, whereas, it was worked out Rs. 38913.68, Rs. 35111.90, Rs. 33692.41 and Rs. 31527.87 on marginal, small, medium and large size of sample farms respectively with an average of Rs. 36099.09 for second cluster. Overall average of cost C was observed Rs. 36541.27 for both the clusters.

Overall, cost of cultivation of potato per



hectare on cost Ai/A, was worked out Rs. 20971.18. Cost B1 Rs. 25639.04, cost B2 Rs. 29289.04, cost C1 Rs. 32891.27 and cost C, Rs. 36541.27, respectively.

It was observed that farm production of potato gross return was more on marginal farms than that of small, medium and large farms in case of both the clusters, because of more use of variable inputs by these farmers as compared to other. On an average, gross returns came to Rs. 97366.04 and Rs. 96564.52 for first and second cluster while overall gross returns came to Rs. 96965.28. Net returns over cost C1 and C2 were calculated Rs. 64182.60 and Rs. 60382.60 for first cluster and Rs. 63965.43 and Rs. 60465.43 for second cluster. Overall average came to Rs. 64074.02 and Rs. 60424.02 per hectare, respectively. The findings were in line with the research results of Singh et al.(1991) and Pandit et al. (2002).

Similarly, family labour income, farm investment income and farm business income were worked out to Rs. 67455.83, Rs. 45093.25 and Rs. 75565.64, respectively in case of first cluster whereas, in case of second cluster, it was Rs. 67896.66, Rs. 44624.99 and Rs. 76422.56, respectively. The overall average was observed Rs. 67676.25, Rs. 44859.12 and Rs. 75994.10, respectively.

Cost of production per quintal of potato was computed to Rs. 176.88 in case of first cluster and Rs. 174.02 for second cluster. Overall cost of production/quintal was computed to Rs. 157.91 and Rs. 175.45, respectively. On an average, input-output ratio on the basis of cost Ai/A2, cost B1, cost B2, cost C1 and cost C2 were worked out to 4.99, 3.73, 3.26, 2.94 and 2.64, respectively for 1st cluster whereas, in case of second cluster, it was worked out 4.80, 3.84, 3.37, 2.98 and 2.69, respectively. On

overall average was 4.65, 3.79, 3.32, 2.96 and 2.67, respectively.

Marketable and Marketed Surplus of Potato

The nature and extent of marketable and marketed surplus of potato per farm is presented in Table 3.

Table 3 revealed that average marketable surplus in cluster first accounted 41.43 quintals i.e. 83.92 per cent of the total production which varied from 72.22, 80.96, 81.55 and 90.61 per cent on marginal, small, medium and large farms, respectively. While in case of second cluster average marketable surplus was worked out 34.46 quintal i.e. 80.91 per cent of the total production, which varied from 72.60, 76.52, 82.23 and 85.15 per cent of total production on marginal, small, medium and large farms, respectively. Overall average of marketable surplus came to 37.95 quintals i.e. 82.53 per cent of the total production.

Average marketed surplus was worked out 40.29 quintal i.e. 81.61 per cent, which varied from 70.06, 77.92, 78.68 and 88.96 per cent of the total production on marginal, small, medium and large farms, respectively on first cluster. While in case of second cluster, average marketed surplus came to 32.83 quintal i.e. 77.08 per cent which varied from 69.62, 70.42, 78.28 and 82.09 per cent of the total production on marginal, small, medium and large farms, respectively. Overall average of marketed surplus was observed to 36.56 quintal per farm i.e. 79.51 per cent of the total production. It can be inferred from the table that marketable and marketed surplus was increasing with the increase of size of sample farms in both the clusters. The findings were in line with the research results of Dahiya&Pandey, (1992).

Table 1 per hectare cost of different inputs used on different size group of potato

Items	1 st cluster					II nd cluster					(Rs.)
	Marginal	Small	Medium	Large	Average	Marginal	Small	Medium	Large	Average	Overall
Family labour	8980.27 (27.12)	6720.87 (20.39)	5070.54 (15.87)	3848.24 (12.20)	7073.23 (21.65)	8796.73 (27.56)	7416.52 (24.00)	6113.97 (20.26)	4432.19 (14.93)	7431.23 (23.92)	7252.23 (22.75)
Hired labour	112.60 (0.34)	1062.26 (3.22)	2228.56 (6.98)	3025.27 (9.59)	1089.90 (3.34)	120.88 (0.38)	1160.15 (3.75)	1748.91 (5.80)	2940.76 (9.91)	1022.52 (3.29)	1056.21 (3.31)
Bullock power	1120.39 (3.38)	845.95 (2.57)	520.17 (1.63)	247.00 (0.78)	836.20 (2.56)	913.17 (2.86)	698.68 (2.26)	602.48 (2.00)	298.85 (1.01)	727.76 (2.34)	781.98 (2.45)
Tractor & Machineries	586.43 (1.77)	1075.64 (3.26)	1198.82 (3.75)	1390.00 (4.41)	920.28 (2.82)	657.54 (2.06)	827.23 (2.68)	1243.57 (4.12)	1334.56 (4.50)	888.40 (2.86)	904.34 (2.84)
Seed cost	13425.38 (40.55)	14105.37 (42.81)	14120.90 (44.20)	14180.28 (44.94)	13817.97 (42.29)	12899.48 (40.41)	12137.21 (39.27)	12116.33 (40.16)	12520.97 (42.18)	12514.90 (40.28)	13166.43 (41.32)
Manure &	2682.36	2837.53	2515.33	2580.40	2680.41	2627.52	2662.17	2410.18	2218.89	2548.37	2614.39

fertilizer	(8.10)	(8.61)	(7.87)	(8.18)	(8.20)	(8.23)	(8.61)	(7.99)	(7.48)	(8.20)	(8.20)
Irrigation charges	1310.81 (3.96)	1312.24 (3.98)	1185.18 (3.71)	1098.17 (3.48)	1263.05 (3.86)	1448.78 (4.54)	1421.57 (4.61)	1218.35 (4.04)	1208.23 (4.07)	1371.36 (4.41)	1317.21 (4.13)
Plant protection	498.38 (1.51)	545.21 (1.65)	634.33 (1.99)	686.83 (2.17)	557.64 (1.71)	368.39 (1.15)	498.51 (1.61)	620.14 (2.06)	594.75 (2.00)	474.70 (1.53)	516.17 (1.62)
Interest on working capital	592.09 (1.79)	653.53 (1.98)	672.10 (2.10)	696.24 (2.21)	634.96 (1.94)	588.14 (1.84)	582.17 (1.88)	598.80 (1.98)	633.51 (2.13)	593.95 (1.91)	614.46 (1.93)
Rental value of owned land	3800.00 (11.48)	3800.00 (11.53)	3800.00 (11.90)	3800.00 (12.04)	3800.00 (11.63)	3500.00 (10.97)	3500.00 (11.33)	3500.00 (11.59)	3500.00 (11.79)	3500.00 (11.26)	3650.00 (11.45)
Total	33108.71	32958.63	31945.93	31552.43	32673.64	31920.63	30904.21	30172.73	29682.71	31073.19	31873.42

(Value in parentheses denote percentage)

Table 2 Measures of per hectare costs and returns of potato

(Rs.)

S.No	Items	Ist cluster					II nd cluster					
		Marginal	Small	Medium	Large	Average	Marginal	Small	Medium	Large	Average	Overall
1.	Cost A1/A2	20328.44	22437.76	23075.39	23904.19	21800.40	19623.90	19987.69	20558.78	21750.52	20141.96	20971.18
2.	Cost B1	262160.97	26582.73	25509.99	25433.97	26110.21	26616.95	24195.38	24078.44	23595.68	25167.86	25639.04
3.	Cost B2	30060.97	30382.73	29309.99	29233.97	29910.21	30116.95	27695.38	27578.44	27095.68	28667.86	29289.04
4.	Cost C1	34241.24	33303.60	30580.53	29282.21	33183.44	35413.68	31611.90	30192.41	28027.87	32599.09	32891.27
5.	Cost C2	39041.24	37103.60	34380.53	33082.21	36983.44	38913.68	35111.90	33692.41	31527.87	36099.09	36541.27
6.	Gross income	98792.00	97394.00	95996.00	94132.00	97366.04	98326.00	96928.00	94598.00	92268.00	96564.52	96965.28
7.	Net return over cost C1	63550.76	64090.40	65415.47	64849.79	64182.60	62912.32	65316.10	64405.59	64240.13	63965.43	64074.02
8.	Net return over cost C2	59750.76	60290.40	61615.47	61049.79	60382.60	59412.32	61816.10	60905.59	60740.13	60465.43	60424.02
9.	Family labour income	68731.03	67011.27	66686.01	64898.03	67455.83	68209.05	69232.62	67019.56	65172.32	67896.66	67676.25

S.No.	Family	Name of the Macrophytes	Habitat
1.	Typhaceae	Typha augustata	Emergent
2.	Potamogetonaceae	Potamogeton natans	Floating
3.	Hydrocharitaceae	Vallisneria spiralis, Hydrilla verticillata	Submerged
4.	Polygonaceae	Polygonum serotinum	Emergent
5.	Trapaceae	Trapa bispinosa	Floating
6.	Pontederiaceae	Eichhornia crassipes	Floating
7.	Ceratophyllaceae	Ceratophyllum demersum	Submerged
8.	Nymphaeaceae	Nelumbo nucifera, Nymphaea stellata	Floating
9.	Azollaceae	Azolla pinnata	Floating
10.	Acanthaceae	Asteracantha longifolia	Emergent
11.	Lentibulariaceae	Utricularia flexuosa	Submerged
12.	Cyperaceae	Cyperus pangarensis	Emergent
13.	Ipomoeaceae	Ipomoea aquatica	Submerged
14.	Plantaginaceae	Limnophylla aquatica	Emergent

Particulars			IstCluster			IInd cluster					
	Marginal	Small	Medium	Large	Average	Marginal	Small	Medium	Large	Average I	Overall
Production	15.26	41.48	62.94	171.17	49.37	15.11	29.99	57.52	148.27	42.59	45.98
Family consumption	2.02	3.56	2.33	4.44	2.77	2.18	2.53	2.77	4.57	2.66	2.72
	(13.24)	(8.58)	(3.70)	(2.59)	(5.61)	(14.43)	(8.44)	(4.82)	(3.08)	(6.25)	(5.91)
Seed	1.51	2.77	6.22	8.50	3.52	1.25	2.92	5.17	14.37	3.96	3.74
	(9.89)	(6.68)	(9.88)	(4.47)	(7.13)	(8.27)	(9.74)	(8.99)	(9.69)	(9.31)	(8.13)
Wages	0.71	1.57	3.06	3.13	1.65	0.71	1.59	2.28	3.08	1.50	1.58
	(4.65)	(3.78)	(4.86)	(1.83)	(3.34)	(4.70)	(5.30)	(3.96)	(2.08)	(3.53)	(3.43)
Losses	0.33	1.26	1.81	2.82	1.14	0.45	1.83	2.27	4.53	1.63	1.39
	(2.16)	(3.04)	(2.88)	(1.65)	(2.31)	(2.98)	(6.10)	(3.95)	(3.06)	(3.83)	(3.02)
Marketable surplus	11.02	33.58	51.33	155.10	41.43	10.97	22.95	47.30	126.25	34.46	37.95
	(72.22)	(80.96)	(81.55)	190.61	(83.92)	(72.60)	(76.52)	(82.23)	(85.15)	(80.91)	(82.53)
Marketed surplus	10.69	32.32	49.52	152.28	40.29	10.52	21.12	45.03	121.72	32.83	36.56
	(70.06)	(77.92)	(78.68)	(88.96)	(81.61)	(69.62)	(70.42)	(78.28)	(82.09)	(77.08)	(79.51)**

(Values in parentheses denote percentage)

****Overall**



CONCLUSION

It may be concluded that Overall, cost of cultivation per hectare was observed Rs. 31873.42, the major share of cost was 41.32 per cent of seed followed by 22.75 per cent family labour and 8.20 per cent manure and fertilizer, respectively.

On an average, gross return, net return, family labour income, farm investment income and farm business income Rs. 98965.25, Rs. 60424.02, Rs. 67676.25 and Rs. 44859.12, respectively. Overall, cost of production per quintal on the basis of cost C1 and C was found Rs. 157.91 and Rs. 175.45, respectively. On an overall average, input-output ratio was found 2.67 per cent. Overall average of marketable surplus came to 37.95 quintal i.e. 82.53 per cent of the total production and overall average of marketed surplus was recorded to 36.56 quintal per farm i.e. 79.51 per cent of the total production.

REFERENCES

1. Neoplasia-Cell Biology Fundamentals and applications, M.L. Gupta and M.L. Jangir., *Agrobios* **2001**.634-652.
2. Medicinal plants may help treat cancer : Experts say – 13 Jan **2004**., Dawn the INTERNET Edition , [http:// DAWN.com](http://DAWN.com) staff reporter.
3. Antioxidants- <http://www.health-caremagic.com/health> Resource.
4. Medicinal plants vs cancer. Henrylito D. Tacio. Sept 19-**2005**. [http:// sunstar.com.ph/e-magazine/index.htm](http://sunstar.com.ph/e-magazine/index.htm).
5. Understanding free radicals and antioxidants. <http://www.healthcheck-system.com/info.htm>.
6. Cancer: a single disease with multitude manifestations? Peter Grandics. *J. of Carcinogenesis*. **2003**,2-9.
7. Can Medicinal Plants Beat Cancer? Keith Stelling, MNIMH, Dip-Phyt alive #222, April **2001**.
8. Ethano medicinal plants for prevention and treatment of tumors. Sharma and Pandey. *International journal of green pharmacy march* **2009**. 2-5.
9. Gaidhani SN, Lavekar GS, Juvekar AS, Sen S, Singh A, Kumari S. In-vitro anticancer activity of standard extracts used in ayurveda. *Phcog Mag* **2009**;5:425-9.
10. Gupta V K, Sharma S K. plants as naturals anti- oxidants. Nat. Pro. Rad. 2006, 326-34. Kaur C, Kapoor H C., Anti- oxidant activity and total phenolic content of some Asian vegetables. *Int. J. Food Sci. Tech.* **2002**; 37;153-61)
11. Chatterji KK. A note on margo-sa oil in the treatment of cancer. *Lancet* **1925**;2:1063-4.
12. Chandra D, Gupta SS. Antiinflammatory and antiarthritic activity of volatile oil of Curcuma



longa (haldi). *Indian J Med Res* **1972**;60:138-42.

13. Hastak KA, Lubri N, Jakhi SD, More C, John A, Ghaisas SD, *et. al.*, Effect of turmeric oil and turmeric oleoresin on cytogenetic damage in patients suffering from oral sub-mucous fibrosis. *Cancer Lett* **1997**; 116:265-69. *urcuma longa* (haldi). *Indian J Med Res* 1972;60: 138-42.
14. Bhide SV, Azuine MA, Lahiri M, Telang NT. Chemoprevention of mammary tumor virus-induced and chemical carcinogen-induced rodent mammary tumors by natural plant products. *Breast Cancer Res Treat* **1994**; 30:233-42.
15. Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics, **2002**. CA: *Cancer J Clin* 2005;55:74-108.
16. Shoeb M. Anticancer agents from medicinal plants. *Bangl J Pharmacol* **2006**;1:35-41.
17. Adebayo AH, Tan NH, Akindahunsi AA, Zeng GZ, Zhang YM. Anticancer and antiradical scavenging activity of *Ageratum conyzoides* L. (Asteraceae). *Phcog Mag* **2010**;6:62-6
18. Late Shringi MS. Personal communication **2002**.
19. Araujo,CC, Leon,LL.Biological activities of *Curcuma longa* L. *Phytother Res.* **2001** Jul; 96 (5):7234-8.
20. Sang Tae Km., Jung Do Kim, Seung Hee Ahn, Young Ik Lee, Young Seok Jeog **2004**. Hepato protective & anti oxidant effects of *Alnus japonica* extracts on acetaminophen induced hepato toxicity in rats.*Phytother Res.* **2004**.Dec 18 (12):971-5.
21. Adebayo AH, Tan NH, Akindahunsi AA, Zeng GZ, Zhang YM. Anticancer and antiradical scavenging activity of *Ageratum conyzoides* L. (Asteraceae). *Phcog Mag* **2010**;6:62-6
22. Newman DJ, Cragg GM. Natural products as sources of new drugs over the last 25years. *J Nat Prod* **2007**;70:461-77.
23. Han HJ, Tan NH, Zeng GZ, Fan JT, Huang HQ, Ji CJ, *et. al.*, Natural inhibitors of DNA topoisomerase I with cytotoxicities. *Chem Biodivers* **2008**;5:1364-8.
24. Shukla Y, Kalra N. Cancer chemoprevention with garlic and its constituents. *Cancer Lett* **2007**; 247:167-81.
25. Adebayo AH, Tan NH, Akindahunsi AA, Zeng GZ, Zhang YM. Anticancer and antiradical scavenging activity of *Ageratum conyzoides* L. (Asteraceae). *Phcog Mag* **2010**;6:62-6.





EFFECT OF PROFENOFOS (INSECTICIDE) ON HEMATOLOGICAL PARAMETERS OF OCCUPATIONALLY EXPOSED WORKERS

Sadhana kesharwani and khushboo sahu

Dept. of Zoology, Govt. M.H. College of Home science &
science for Woman, Jabalpur (M.P.)

Email : sadhanakesharwani.23@gmail.com, khussahu28@gmail.com

(Date of Receipt :04-01-2015 ;

Date of Acceptance for Publication : 31-01-2015)

Abstract

Profenofos (O-4-bromo-2chlorophenyl O-ethyl S-propyl phosphorothioate) are chemical substances used in agricultural production to protect crops against insects. They help to achieve better quality and quantity of crops; however, they also are capable of causing occupational diseases in farmers and workers. During the spraying time we collected the blood sample of workers during month of November(2014) to January(2015) from different villages of Seoni district (M.P.). In the present study, total 26 blood samples were collected out of which 21 persons were exposed to profenofos insecticide. They were compared with 5 control persons. Different blood parameters and liver enzymatic test were determined in comparison to control subjects. The results indicate some alterations in blood indices and liver enzymatic test in all the exposed workers. Low MCV, MCH, PCV was noted in almost all the exposed worker. High lymphocyte, eosinophil, neutrophil, MCV, MCH was noted in some workers. Only two workers found with high liver bilirubin and SGPT. Survey of exposure indicated that prolonged profenofos exposure affected the health of Workers and produced dermatological, respiratory and other clinical disorders.

Key Words: Profenofos, Blood Parameters, Human Health.

Pages: 07

References: 12

INTRODUCTION

Profenofos is a natural toxic organophosphorous insecticide which kill insects in field (Ismail *et. al.*, 2010). The use of organophosphorous insecticides has increased food production with increasing population in many parts of the world. (Al-Shinnawy, 2008). Exposure to profenofos

in the recent years has increased manifold due to their intense application and as such the risk of occupational hazards have also multiplied tremendously. According to WHO (2009), approximately 25 million profenofos poisoning cases occur annually among agricultural workers in developing countries.



There is doubt that some persistent insecticide accumulate in various biological system at level much higher than those in their surroundings. There are 3 major routes of transport of this insecticide within the body that may be through leaching water, aerial contamination and accidental or professional exposure. Agricultural workers are exposed to profenofos primarily through mixing of chemicals, loading into dispensers, application, clean up and disposal of empty chemical containers (Wolf *et. al.*, 1967).

Profenofos affects the blood parameters of occupationally exposed workers. Correlation of different blood parameters with the harmful effects of profenofos especially in the case of total leukocytes count (TLC), red cell count (RBC), WBC, hemoglobin content (Hb%), hematocrit/packed cell volume (PCV), differential leukocyte counts (DLC), MCV, MCH, MCHC and platelet count has been reported by many researches (Srivastava *et. al.*, 1991, Mythilal, 2000, Lee *et. al.*, 2002, Corsini *et. al.*, 2005). Much work has been done in many countries on profenofos blood parameter correlation. Profenofos is largely known to causes different symptoms in liver and kidney. Profenofos causes different symptoms of toxicity and biochemical changes in the enzyme activity of the liver like- SGOT (Serum Glutamic Oxaloacetate Transaminase) , SGPT (Serum Glutamic Pyruvate Transaminase) (Reitmann *et.al.*,1957). Profenofos are also affect liver Bilirubin , Biliviridin as well as Acid and Alkaline phosphateses level.

Most occupational exposures to this insecticide occur from skin absorption and inhalation as the sprayers are from poor family and not in a habit to use face

masks. Farm workers, traders and people employed in the manufacture of pesticides are especially at risk of pesticides exposure. Such risks are particularly pronounced in developing nations, where hazards are commonly less well understood healthwise and safety regulations are less stringent or poorly enforced. Accidental oral contamination can occur when farmers eat, drink or smoke while spraying or do so shortly after spraying without washing their hands.

MATERIAL AND METHODS

Total 26 Blood samples taken during spraying time from November(2014), to January(2015). Out of which 21 persons were exposed to profenofos and 5 normal control persons(not exposed to profenofos) taken for the determination of profenofos residues and different blood parameters and liver enzymatic parameters.

The subjects were selected from Seoni District. Profenofos sprayers were selected from agricultural workers/farmers in different age groups. Agricultural workers and traders exposed to profenofos were interviewed and their details with regard to age, smoking habit, sex and duration of exposure were reported complete questionnaire in which personal data, working activity, type and duration of contact with profenofos, smoking habits, precautions taken during exposure to profenofos, and any physiological problem after exposure were recorded.

The following hematological parameters were studied to correlate any possible relation with the other studied parameters- R.B.C.count, W.B.C count, MCV,MCH,MCHC, differential leukocyte count(DLC), Hemoglobin percentage, hematocrit value/PCV. Enzymatic test

like- SGOT(Serum Glutamic Oxaloacetate Transaminase), SGPT(Serum Glutamic Pyruvate Transaminase), Billirubin and Acid and Alkaline phosphateses level. These blood parameters were studied using standard hematological procedures.

RESULTS AND DISCUSSION

The present work showed that the blood samples of 21 exposed sprayers, 8 peo-

ples were found affected. These people showed low level of MCV, MCH, neutrophill and eosinophill, out of which 4 persons were prolonged smoker. Other 4 persons showed high level of MCHC, neutrophill, eosinophil. Only 2-3 persons were found with low level of lymphocyte and htc/pcv . 1-2 persons found with high level of Hct and lymphocyte. as well as 2 persons were reported with high level of liver bilirubin and SGPT.

Table-1(a): Determination of blood parameters in the Profenofos exposed workers/traders-

Sample no.	Blood Parameters												
	Hb	TLC	N	E	B	L	M	HCT	RBC	MCV	MCHC	MCH	PLT
1	12.6	5.9	60	04	00	34	02	37.3	4.98	75.1*	33.7	25.3*	167
2	14.1	8.8	46*	07*	00	45*	02	39.0	5.80*	67.4*	36.1*	24.3*	229
3	12.4	7.8	54	07*	00	36	03	35.2*	5.15	68.5*	35.2	24.0*	137
4	13.5	7.2	60	05	00	33	02	39.0	4.74	82.4	34.6	28.4	155
5	12.1	6.3	60	05	00	33	02	34.4*	4.80	71.7*	35.1	25.2*	170
6	12.4	6.4	69	04	00	25	02	36.2	5.4	67.2*	34.2	22.9*	118
7	14.8	5.7	56	05	00	37	02	40.3	4.25	94.9	36.7*	4.8*	169
8	13.4	7.6	67	05	00	26	02	37.2	4.11	72.9*	36.0	26.2	131
9	12.0	9.4	60	07*	00	30	03	34.0*	4.64	73.4*	35.2	25.8*	185
10	12.8	6.5	44	05	00	49*	02	36.1	4.25	85.1	35.4	30.1	112
11	13.7	8.4	74	04	00	20	02	39.5	5.68*	69.7*	34.6	24.1*	128
12	13.7	5.5	52	07*	00	38	03	39.2	4.57	85.9	34.9	29.9	134
13	12.0	9.6	82*	02	00	15*	01*	36.8	5.14	71.6*	32.6	23.3*	207
14	14.7	8.8	63	05	00	30	02	42.3	4.98	85.1	34.7	29.5	151
15	13.7	8.9	57	07*	00	34	02	37.6	5.75*	65.4*	36.4	28.3*	150
16	13.2	9.7	77*	07*	00	14.1*	02	35.8*	4.5	79.5*	36.8*	29.2	345
17	12.1	6.4	63	05	00	34	04	34.7*	4.70	72.3*	35.1	25.4	181
18	12.1	6.3	55	05	00	38	02	38.8	3.94	98.5	38.9*	38.3*	198
19	15.4	6.6	62	03	00	33	02	39.4	4.69	84.2	39.0*	32.8*	173
20	13.4	7.6	63	07*	00	18*	02	38.2	4.62	99.3*	31.5	36.5*	185
21	15.9	6.3	64	05	00	29	02	42.0	4.46	94.2	37.8*	35.6*	231

Table 1(b):

Sample No.	Liver enzymatic parameters			
	SGOT	SGPT	Bilirubin	Alkaline phosphate
1	35.07	40.06*	1.44*	110.02
2	29.42	31.07	0.94	89.74
3	28.06	30.05	0.72	90.42
4	20.06	24.00	0.36	80.14
5	20.06	34.08	1.08*	95.17
6	18.06	24.08	0.72	92.10
8	31.09	27.49	0.79	118.04
9	22.90	14.92	1.44*	84.70
10	19.85	25.02	0.28	150.03
11	30.49	37.05*	2.52*	85.43
12	17.08	21.30	0.41	227.01
13	37.02*	16.10	0.91	113.71
14	12.70	26.03	0.80	215.68
15	13.90	15.06	0.69	116.17
16	19.42	20.23	0.51	91.06
17	19.33	18.84	0.61	69.07
18	14.87	20.91	0.48	79.09
19	34.01	30.02	0.92	123.22
20	27.05	31.89	0.86	132.04
21	26.02	17.05	0.93	149.03

Note- In this tables the affected blood parameters were showed by ‘*’

In the control group only 2 person were found . out of which one affected from low level of MCV and second affected from the low level of MCH. They are not affected regarding other parameters.

Table 2(a): Determination of the blood parameters in the blood of normal (control) workers-

Sample no.	Blood Parameters												
	Hb	TLC	N	E	B	L	M	Hct	RBC	MCV	MCHC	MCH	PLT
1	13.4	6.6	69	5	0	33	2	39.1	4.26	84.3	33.7	25.2*	197
2	12.6	7.1	52	5	0	28	2	37.3	3.80	91.1	32.8	29.5	196
3	12.8	5.6	57	6	0	27	4	46.4	5.21	97.5	34.1	29.1	191
4	14.2	4.9	61	4	0	31	2	40.3	4.90	78.4*	35.2	27.2	163
5	13.7	6.3	64	3	0	30	3	45.5	3.75	86.3	33.6	30.3	159

Table 2(b):

Sample No.	Liver enzymatic parameters			
	SGOT	SGPT	Bilirubin	Alkaline phosphate
1	31.80	24.50	0.56	116.10
2	19.42	16.20	0.89	122.15
3	27.05	30.01	0.77	99.08
4	35.40	17.05	0.71	91.03
5	14.83	19.04	0.69	113.60

Normal values: Hb = hemoglobin [11.0-16.0 g/dl], TLC=total leucocyte count[4000-5000/cumm], N=Neutrophill[50-75%], E=Eosinophill[1-5%], B= Basophill[0-1%], L=Lymphocyte[20-40], M=Monocyte[2-5%], Hct=Hematocrit[36-48%], RBC = Red blood cell count[3.5-5.5mil/cmm] , MCV=Mean corpuscular volume[80-99fl], MCHC=Mean corpuscular hemoglobin concentration[32-36gm/dl], MCH= Mean corpuscular hemoglobin[26-32pg], PLT=Platelet count[100-400 Thsd /cmm].

Azmi et.al., (2009), reported some alterations in blood indices in all the insecticide exposed persons, and only two persons from two different stations were affected severely and therefore their Hb, MCV, MCHC, TLC, monocyte and neutrophil counts increased significantly. High lymphocyte count was noted in almost all the exposed persons. Platelets count was also found high in few persons but Hb, MCV, MCH, TLC, RBC and neutrophil counts significantly decreased in such persons. Exposure of multiple pesticides for prolonged period has also affected the health of exposed persons and produced dermatological, hepatic, nephritic, respiratory and other clinical disorders reflecting the toxic effects of pesticides.

Lu et.al., (2007), reported abnormal eosinophil count in insecticide sprayers. Increment of blood eosinophill level was observed in subjects that suffering from cough, wheezing, eye and nasal irritation and other allergic symptoms like itching, sneezing, skin rashes and breathlessness. **Kossmann et.al., (1993)**, observed high lymphocyte count in almost all the exposed persons from different farm stations. Platelet count was found low in almost all the exposed cases. Only few per-

sons showed high platelet counts in some stations due to the frequent and direct exposure of profenofos insecticides during their working time but their Hb, Hct, MCV, MCH, RBC and neutrophil counts were found in low levels in such persons.

REFERENCES

1. AL-Shinnawy, MSA. (2008). Effects of orally fed diazinon on some biochemical parameters of male albino rats. *The Egyptian journal of hospital medicine*, 33, 559-568.
2. Azmi, M.A., S.N.H. Naqvi and M.A. Azmi.(2009). pesticide residues in the blood of rural population from gadap, Karachi and related health hazards. *J. Environ.Biol. India*, 30(5), 747-756.
3. Corsini, F.S., Birindelli, S. Fustioni, G.De Paschale, T. Mammone, S. Visentin, G.L. Galli, M. Marinvovich and colosio. (2005): immunomodulatory effects of the fungicide mancozeb in Agricultural workers. *Toxicol. Appl. Pharmacol.*, 208, 178-185.
4. Ismail AA, Rohlman Ds, Abdel RGM (2010). clinical and Biochemical parameters of children and adolescents applying pesticides. *Int. J. Occup environ Med.* 3:132-143.
5. Kossmann, S.J., Kosmider and B. Konieczny (1993): cytoenzymatic examinations of peripheral blood granulocytes in overhaul workers of the chemical plant "Organika-Jaworzno" at Jaworzno. *Med. Pr.*, 44,35-39.
6. Lee, E.Y., Y.K. Hwang, J.O. vang and S.V. Hong (2002); predictors of survival after acute piragua poisoning. *Toxicol.*



Ind. Hlth., 18, 201-206 .

7. Lu, J.L.(2007). Acute pesticide poisoning among cut-flower farmers. *J. Envir. Health.*,70(2), 38-43
8. Mythilai, A. (2000): Endosulfan Poisoning – A clinical profile. *J. Assoc. physicians. India*, 48, 323-325 .
9. Reitmann, S., Frankel,S.(1957). For the determination of SGOT and SGPT Activity in serum. *Amer. J. Clin. Path.*,28: 56.
10. Srivastava, A.K., B.N. Gupta, A.K. Mathur, N. Mathur, P.N. Mahendra and R.S. Bharti (1991): The clinical and biochemical study of pesticide sprayers. *Hum. Exp. Toxicol.*, 10, 279-283 .
11. W.H.O., (1998). The WHO recommended classification of pesticides by hazard and guideline to the classification 1988-99 WHO/VBC/88.953, world health organization, Geneva.
12. Wolf, H.R. Durham, W.F. and Armstrong J.F.,(1967). Exposure of workers to pesticides. *Archives of Environmental Health.*, 14, 622-33.





DIVERSITY OF MACROPHYTE COMMUNITY OF A LENTIC WATER BODY OF JABALPUR

Dr. Reeta Solanki

Deptt. of Zoology & Biotechnology,
Govt. M.H. College of H.Sc. & Sci.,
Jabalpur (M.P.)

Email : solanki1111@rediffmail.com

(Date of Receipt :14-01-2015 ;

Date of Acceptance for Publication : 11-02-2015)

Abstract

The study was conducted to observe the diversity of macrophyte community of Robertson Lake of Jabalpur. The macrophyte community comprised of 16 species belonging to 14 families & showed seasonal fluctuation.

Key words :- Diversity, Macrophytes, Aquatic Macroscopic Plants.

Pages: 3

References: 8

INTRODUCTION

Macrophytes, the aquatic macroscopic plants also known as hydrophytes. Confine themselves to the littoral zone of the lake & contribute significantly to the total primary production.

The hydrophytes exhibit poor mechanical, conductive absorbing & protective tissues & possess air spaces in the tissues. They play an important role in the productivity & nutrient status. They provide an enormously large surface area for the growth of microflora & fauna and help in anchoring the fish spawn but sometimes they hinder the growth of aquatic animals, particularly fishes. They utilize mineral elements from the bottom sediments and hereby affecting nutrient

level of aquatic ecosystem (Deka *et. al.*, 2012).

Studies on different aspects of macrophytes of fresh water ecosystem were carried by Odum (1959), Mishra & Tripathi (2004), Sobha & Harilal (2005), Mathur & Mathur (2006) & Kiran *et. al.*, (2007).

The importance of macrophytic studies in fresh water ecosystem was recognized quite late as compared to the studies on physico chemical and phytoplanktonic parameters.

STUDY AREA

Robertson lake is one of the largest lake

left behind by Gondwana Regime. The lake spreads over the area of 67 hectare being situated in a remote suburban area of the city. This lake is in a better condition as compared to most of the other ponds in the city, which have fallen prey to the greed of land Mafia or are being swallowed by the slums erupting like cancer-

ous sores on the bank of the pond.

MATERIAL AND METHOD

For the present study five sites were chosen (i.e. East, West, North, South & Central). The macrophytic species were collected & identified with the help of Burche (1991) & Cook (1996).

RESULT & DISCUSSION

The list of macrophytes of the Robertson lake is presented in Table No. 1.

S.No.	Family	Name of the Macrophytes	Habitat
1.	Typhaceae	Typha augustata	Emergent
2.	Potamogetonaceae	Potamogeton natans	Floating
3.	Hydrocharitaceae	Vallisneria spiralis, Hydrilla verticillata	Submerged
4.	Polygonaceae	Polygonum serotinum	Emergent
5.	Trapaceae	Trapa bispinosa	Floating
6.	Pontederiaceae	Eichhornia crassipes	Floating
7.	Ceratophyllaceae	Ceratophyllum demersum	Submerged
8.	Nymphaeaceae	Nelumbo nucifera, Nymphaea stellata	Floating
9.	Azollaceae	Azolla pinnata	Floating
10.	Acanthaceae	Asteracantha longifolia	Emergent
11.	Lentibulariaceae	Utricularia flexuosa	Submerged
12.	Cyperaceae	Cyperus pangareii	Emergent
13.	Ipomoeaceae	Ipomoea aquatica	Submerged
14.	Plantaginaceae	Limnophylla aquatica	Emergent



The macrophytic communities comprises of a 14 families, represented by 16 species.

Out of families 5 sps were emergent 7 sps floating 5 sps. were from submerged habitat.

During the present study, free floating herb *Eichhornia crassipes* was observed as most dominant macrophyte, on the banks & towards the inlets.

Other major macrophytes observed during studies were water spinach (*Ipomoea aquatica*) & *Nelumbo nucifera*.

Most of the macrophytic communities were perennial but their occurrence was found decreased in rainy season & found evident during post monsoon periods.

This study reveals that the large part of the lake is covered by macrophytes. In shallow areas the spreading of macrophyte reduces areas available to the aquatic animals & obstruct their mobility.

REFERENCE

1. Bursche, E.M. (1991). A Handbook of water plants, Oriental Enterprise, Dehradun.
2. Cook, C.D.K. (1996). Aquatic & Wetland plants of India, Oxford University Press, New Delhi, India.
3. Deka, R.M., Bauah, B.K., Kalita, J. (2012). Study on diversity & quantitative characteristics of macrophytes of Kapla Beel, A tropical flood plain wet land in Assam. *Poll. Res.* 31 (4) : 639-642.
4. Kiran, B.R., Hamsa A. & Puttaiah E.T. (2007). Aquatic & Marshy plants and their economic importance in Bhadra reservoir project region, Karnataka. *Nature Environment & Pollution Technology*, 6 (3) : 429-432.
5. Mathur, S.M. & Mathur P. (2006). Aquatic weeds & its impact on environment with special reference to water hyacinth. *Ecol. Env. Cons.* 12 (3) : 535-542.
6. Mishra, B.P. & Tripathi, B.D. (2004). Distribution of macrophytes & phytosociology of *Hydrilla Verticellata* Casp. & *Lemna minor* Linn. in lotic & lentic aquatic ecosystems. *Ecol. Env. & Cons.* 10 (1) : 37-41.
7. Odum, E.P. (1959). Fundamentals of Ecology, II edition, W.B. Saunders Company. Philadelphia.
8. Sobba, V. & Harilal, C.C. (2005). Influence of changing water quality of *Eichhornia Crassipes* (Mart.) solms. in the uptake patterns of various minerals & nutrients. *Ecol. Env. & Cons.* 12 (1) : 1-6.







Prof. T.R.C Sinha a true lover of science and always concerned about the environment. It was his dream to publish scientific journals and he started this project with the same enthusiasm as he had done for every project he undertook. We are guided by the vision of Prof. Sinha and endeavour to make his dream of promoting science and help young scientists, publish their articles and encourage them in their research.

Manisha Verma



Submit your manuscripts

@ manisha_npp @ yahoo.com



manishanpp

Academic and Research Publications





**International Journal
Of
Agricultural Science & Technology**
(Bi-Annual Journal)
2015

SUBSCRIPTION ORDER FORM

Name: _____

Address: _____

Pin: _____
Tel: _____ Fax: _____
E-mail: _____

MODE OF PAYMENT

Payment should be made online. The amount is to be credit in the Bank The details of the bank are as follows:-

Name of the Account holder : **Academic And Research Publications**

Name of the bank : **PUNJAB NATIONAL BANK, VAISHALI BRANCH, GHAZIABAD,**
IFSC: PUNB0405300

Current Account No. 4053002100011106

New

☐

Renewal

☐

SignatureDate

Published by

Academic And Research Publications

Office: 22, Gaur Galaxy, Plot No 5, Sec-5, Vaishali , Ghaziabad (U.P.) - 201010 (INDIA)

National Journal of Comparative Law

Brochure

www.manishanpp.com





SUBSCRIPTION RATES

INDIVIDUAL COPY for CO-AUTHORS Only

Inland	Foreign
Rs. 500/-	US\$ 40
ANNUAL SUBSCRIPTION	
(2 Issues Per Year)	

Individuals	Foreign
Rs. 800/-	US\$ 80

Institutions	Foreign
Rs.1500/-	US\$ 100

ADVERTISEMENT RATES



Full Page

Colour	:	Rs. 35,000.00
Black/White	:	Rs. 25,000.00

Half Page

Colour	:	Rs. 25,000.00
Black/White	:	Rs. 15,000.00

**(Four weeks to be allowed after
submission of advertising material)**



Copyrights@www.manishanpp.com 2015- All rights reserved

Submit your manuscripts

@ manisha_npp@yahoo.com, arp@manishanpp.com
www.manishanpp.com

Contact Information

For Subscription orders may be contact directly to publisher (address given below) using Journal Subscription Form or contact your preferred subscription agent. All orders should be accompanied by payment. Please make checks payable to (“**Academic And Research Publications**”). Mail order with payment to:

Subscription Manager :

Academic And Research Publications

Office: 22, Gaur Galaxy,
Plot No 5, Sec-5, Vaishali ,
Ghaziabad (U.P.) - 201010 (INDIA)

Email: manisha_npp@yahoo.com

Phone No.: 0120-4124773

For more information or assistance to place an order, please feel welcome to contact us again at

manisha_npp@yahoo.com

Mode of Payment

Subscriptions are payable in advance and all rates include postage and taxes. Subscribers are requested to send payment with their order whenever possible. Issues will only be sent on receipt of payment. Subscriptions are entered on an annual basis – January to December – and paid in Rs or US Dollars. For payments in Euros and Sterling, please calculate at the prevailing exchange rate.

Payment can be made by Bank (Wire) Transfer, Cheque, International Money Orders. All payments should be made to “**Academic And Research Publications**”.



To order by telephone, please call us at
+0120-4 124 773, 094 15 155 631,
093 103 43 504



JOURNAL SUBSCRIPTION ORDER FORM FOR 2015

Name: _____ Dept. _____
Address: _____
Tel: _____ Mob. _____
E-mail: _____ Pin: _____



Check appropriate boxes (Annual Subscription Price in 1500 Only for *Institution*)



National Journal of Comparative Law(NJCL) ISSN : 2393-9338



International Journal of App. Environmental Science & Technology : 2321-8223



International Journal of Agricultural Science And Technology : ISSN 2319-880X



International Journal of Aquaticscience And Technology : ISSN 2320-6772



International Journal of Nanoscience And Technology : ISSN 2319-8796



International Journal of Geoscience And Technology : ISSN 2321-2144

MODE OF PAYMENT

Payment should be made online. The amount is to be credit in the Bank The details of the bank are as follows:-

Name of the Account holder : Academic And Research Publications

Name of the bank : PUNJAB NATIONAL BANK, VAISHALI BRANCH,
GHAZIABAD,

IFSC: PUNB0405300

Current Account No. 4053002100011106

New

☐

Renewal

☐

SUBSCRIBER TYPE: (Check one) Library

☐

/Institution

☐

/ Personal

☐

Signature

..... Date

Published by

Academic And Research Publications

Office: 22, Gaur Galaxy, Plot No 5, Sec-5, Vaishali, Ghaziabad (U.P.) - 201010 (INDIA)

To order by telephone, please call us at +0120-4124773

Sample Issue: To access free online sample issue of our journals, please send e-mail to manisha_npp@yahoo.com



Dear Reader,

This journal is a collaborative effort which aims to increase awareness and understanding of scientific research, This is a peer reviewed biannual journal which gives a platform to fresh ideas and past experience about developements in India and outside. The journal has Open Access to its content and makes knowledge available to the public in general and research scholars in specific. Through this journal we welcome greater global exchange of knowledge, ideas among intellectuals, researchers and scholars. The journal offers an open forum for discussion. I will like to appreciate the efforts of all authors, co-authors, reviewers and member of Editorial Board who have contributed immencely towards this journal.

ARP. Give your IDEAS an Expression
www.manishanpp.com

Manisha Verma



Copyright: Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, or thesis) that it is not under consideration for publication elsewhere; that if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher.

NOTES



NOTES



Guidelines to Authors

International Journal of Agricultural Science & Technology

This journal is published by the **Academic And Research Publications New Delhi**, twice in a year. The emphasis is to involve a large community of scientists specially JUNIOR SCIENTISTS and scholars from India and abroad in developing a framework of discussion and debate on conservation and sustainable development. Research articles, Review articles Book reviews, Interviews, Short communication, Letters to the Editor, Case reports and News items related to the subject are accepted for publication. The PUBLISHER is aiming to publish your article as rapidly as possible provide a rapid publishing platform, each and every point is vital to reduce the editorial work flow. Authors are requested to extend their maximum cooperation for speedy actions to be taken at this end. Hence they are requested to check the guidelines thoroughly before submitting their article(s) and check that all the required informations are mentioned and strictly according to the guide lines.

Benefit to Authors

Following benefits shall be available to the authors, if your article is published in the journals of this PUBLISHER. Authors can use their published articles in any way they like for the dissipation of knowledge free of charge. However it cannot be used for any commercial use without prior permission. Authors can have access on all articles published by this Publisher. Wide publicity and reach by getting indexed in many free indexing services.

Submission of Article(s)

All manuscripts are to be submitted in English, typed double-spaced throughout the text. It is requested that manuscripts be sent by e-mail only. Authors will be requested to substantiate the need if it exceeds the maximum number of pages.

All manuscript should be submitted to the Publication Editor at manisha_npp@yahoo.com directly.

The Research Article should be in the following order

In the 1st Part:

1) Title 2) Name of the organisation where work was carried out 3) Name of the Author with whom correspondence is to be made along with his/her Mob.No. 4) Date of Submission 5) No. of Tables 6) No. of Photographs 7) Numbers of Graphs 8) No. of References 9) Name and addresses of at least Three referees along with their Email I/Ds and Mob Nos.

In the 2nd Part :

1) Title of The Article 2) Name and addresses of all the Author(s) along with their Email I/Ds and Mob.Nos. 3) Abstract 4) Introduction 5) Materials and methods 6) Results 7) Discussions (if any) 8) Acknowledgements 10) References (Under References heading, Name of the Journals should be in bold letters) and (11) Name of The journal.

Tables

to be included should have a heading, giving the substance, and should be typed double-spaced on separate sheets. They should also be numbered in serial order. Figures either drawn manually or by computer should be in black ink and the lettering on them should be large enough to stand reduction. Photographs in colour should have sharp contrast. Legends for figures and plates should be typed in numerical order on separate sheets, one for figures and one for plates.

References

The literature cited should list the author's name, year of publication, title of the paper, and the Journal titles (bold letters) which should be cited in full (no abbreviation) with volume number and page numbers, as indicated below:

A. For articles in a Journal:-Walsh, J.E. (2008) Climate of the Arctic Marine Environment. Ecological Applications. 18. pp. 3-22.

B. For Books:-Ward, D.R. (2002) Water Wars: drought, floods, folly and politics of thirst: River head Books. New York. p. 12.

C. Chapter in a book:-Andrews, T.J., Cough, B.F. and Muller, G.J. (1984). Photosynthetic gas exchange properties and carbon isotope ratios of some mangroves in North Queensland. In: H.J. Teas (Ed.), Physiology and Management of Mangroves. W. Junk. The Hague. pp. 15-23.

From website:-National Oceans and Atmospheric Administration (NOAA). 1995. Regional Perspectives: Indian Ocean. www.ncdc.noaa.gov/paleo/outreach/coral/sor/sor_indian.html, accessed on July 13, 2008.

While giving reference of more than two authors in the text, after the name of the first author, *et.al.*, should be used, followed by the year of publication.

PRINTING CHARGE

The articles are not printed in gratis. A nominal charge of Rs 500=00 per printed page is charged and is printed in B/W colour. For any coloured page, is separately charged. Charges for the same is quoted on request. Five Reprints and a copy of Journal is provided to the Corresponding author. Extra copies of the Journal for coauthor only are supplied on charge @ Rs 400-00 per copy and shall be sent along with the copy of the Correspondent Author.

REQUEST

Authors are requested to keep a copy of the Mss. till it is published in the Journal.

REMITTANCE

All remittances are to be on line credited in the account of **Academic And Research Publications** in the **PUNJAB NATIONAL BANK, VAISHALI BRANCH, GHAZIABAD, IFSC: PUNB0405300**

Current Account No. 4053002100011106 and inform the at manisha_npp@yahoo.com accordingly.

CONTACT US

For quick reply, please note the address and contact them directly by Post or email:-a) For publication of your article, Acceptance letter, Invoice, sending of Cheques/Drafts, for sending Review Reports, Status Report about your article, and all other queries related to your articles, should be sent directly to the Editor-in-Chief, whose address is as follows:

Prof. Manik Sinha-THE EDITOR-IN-CHIEF, H.Office: EC 41, Maya Enclave, New Delhi -110064

Email: manik.sinha2@gmail.com, manisha_npp@yahoo.com. www.manishanpp.com.

Contents

Reviewed Articles

1. Characteristic of Fog, Foggy Weather and its Impact on Agriculture. 01

Anil Kumar, Raj Singh and S. Singh

2. Economical Analysis Of Potato Production: Nature And Extent Of Marketable & Marketed Surplus 05

Pramod Kumar, R. P. Kaushal, U. A. Siddiqui and Daya Ram

3. Effect Of Profenofos (Insecticide) on Hematological Parameters of Occupationally Exposed Workers 16

Sadhana Kesharwani And Khushboo Sahu

4. Diversity Of Macrophyte Community Of A Lentic Water Body Of Jabalpur 23


Dr. Reeta Solanki

An International Refereed Journal



ARP. Give your IDEAS an Expression

www.manishanpp.com



**ACADEMIC AND RESEARCH
PUBLICATIONS**

**H. Office: EC 41, Maya Enclave,
New Delhi -110064**

Email : manik.sinha2@gmail.com
manisha_npp@yahoo.com

Join us on **facebook**



ARP